


Invariant Large Scale Structure of Axial Diffraction Patterns

Challenge:

- Simple patterns observed in axial plots of intensity.
- No analytic description of patterns known.
- Find (simple) description of patterns.


Background

- Debye scalar wave approximation for focused wave.
- Helmholtz equation and Fourier transform (FT).
- Axial plane is a SLICE \Rightarrow FT is projection of sphere cap.



Observation

- FT of enhanced PSF has sparse curved line structure



- But the analytic FT of (log) enhanced PSF is intractable.
- Intensity has inverse square drop-off with distance.
- Why not use r^2 enhancement instead? Then FT relation is Laplacian:


$$\text{PSF} \Leftrightarrow \text{OTF}$$

$$p(x, z) \Leftrightarrow P(u, w)$$

$$-(2\pi r)^2 p(x, z) \Leftrightarrow \nabla^2 P(u, w)$$

- Laplacian picks out edges and discontinuities of OTF.

Enhanced Intensity PSF and Corresponding OTF



- Enhancement has interesting feature along axis (horizontal).
- Corresponding ∇^2 OTF has vertical lines visible.
- Not really interested in the component along the axis (which is generated by the aperture edge).
- Large scale structure now apparent. Look like overlapping parabolas.
- FT of this structure consists of arcs of circles with dipole-like line cross section.

Initial model of structure

- Consider curvilinear cosine in polar coordinates:

$$p(x, z) = a_1(r)a_2(\theta)\cos[2\pi r f(\theta)]$$


$$x = r \sin(\theta), \quad z = r \sin(\theta)$$

- Particular solutions known (see Amidor for more). Eccentric parabolic cosine:


$$f(\theta) = 1 + \cos(\theta + \beta)$$

$$a_1 = a_2 = 1$$

Eccentric Parabolic Cosine




FT Magnitude




- Analytic solution is two osculating circles with half-order derivative profiles.

Synthesis of simple function with a similar Fourier spectrum

- Modify symmetry of parabolic cosine with shadow edge:




- Add reflected function to fill in the other Fourier quadrants



- Synthesis complete, now compare with computed intensities.

Projection-Slice




- Projection at θ (Radon) \Leftrightarrow Slice of intensity at θ

$$s^{\pm} = \lambda^{-1} [1 - \cos(\alpha \pm \theta)]$$


- Adding mirror image loci gives enhanced intensity composed of overlapping cosine patterns:

$$r^2 I(x, y) \approx \text{const.} [\cos(2\pi r s^+) + \cos(2\pi r s^-)]$$

Intensity loci for 0.5 NA System



Diffraction calculation vs Synthesis



References

- 1 C. J. R. Sheppard, and M. Hrynevych,
"Diffraction by a circular aperture: a generalization of Fresnel diffraction theory.,"
JOSA,A **9**, (2), 274-281, (1992).
- 2 C. W. McCutchen,
"Generalized aperture and the three-dimensional diffraction image,"
JOSA,A **54**, 240-244, (1964)
- 3 A. W. Lohmann,
"Three-dimensional properties of wave-fields,"
Optik **51**, (2), 105-117, (1978).
- 4 N. Streibl,
"Fundamental restrictions for 3-D light distributions,"
Optik **66**, (4), 341-354, (1984).
- 5 C. J. R. Sheppard, M. Gu, Y. Kawata, and S. Kawata,
"Three-Dimensional transfer functions for high aperture systems obeying the sine condition,"
JOSA,A **11**, 593-598, (1994).
- 6 C. J. R. Sheppard, and K. G. Larkin,
"Vectorial pupil functions and vectorial transfer functions,"
Optik **107**, (2), 79-87, (1997).
- 7 K. G. Larkin, and C. J. R. Sheppard,
"Direct method for the phase retrieval from the intensity of cylindrical wavefronts,"
JOSA,A **16**, (7), 1838-1844, (1999).
- 8 Z. Jaroszewicz, A. Kolodziejczyk, D. Mouriz, and S. Bara,
"Analytic design of computer-generated Fourier-transform holograms for plane curves reconstruction.,"
JOSA,A **8**, (3), 559-565, (1991).
- 9 R. N. Bracewell,
Two-Dimensional Imaging,
Prentice Hall, Englewood Cliffs, New Jersey, 1995.
- 10 I. Amidor,
"Fourier spectrum of curvilinear gratings of the second order,"
JOSA,A **15**, (4), 900-913, (1998).